

AD-A266 010



DTIC
S ELECTE D
JUN 22 1993
C

2

AD

TECHNICAL REPORT ARCCB-TR-93016

GENERALIZED MANDELBROT RULE FOR FRACTAL SECTIONS

L.V. MEISEL

APRIL 1993



US ARMY ARMAMENT RESEARCH,
DEVELOPMENT AND ENGINEERING CENTER
CLOSE COMBAT ARMAMENTS CENTER
BENÉT LABORATORIES
WATERVLIET, N.Y. 12189-4050



APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

93 6 21 05

93-13951



DISCLAIMER

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

The use of trade name(s) and/or manufacturer(s) does not constitute an official indorsement or approval.

DESTRUCTION NOTICE

For classified documents, follow the procedures in DoD 5200.22-M, Industrial Security Manual, Section II-19 or DoD 5200.1-R, Information Security Program Regulation, Chapter IX.

For unclassified, limited documents, destroy by any method that will prevent disclosure of contents or reconstruction of the document.

For unclassified, unlimited documents, destroy when the report is no longer needed. Do not return it to the originator.

REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE April 1993	3. REPORT TYPE AND DATES COVERED Final	
4. TITLE AND SUBTITLE GENERALIZED MANDELBROT RULE FOR FRACTAL SECTIONS			5. FUNDING NUMBERS AMCMS: 616224H1800 PRON: 1A12186EA11A	
6. AUTHOR(S) L.V. Meisel				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army ARDEC Benet Laboratories, SMCAR-CCB-TL Watervliet, NY 12189-4050			8. PERFORMING ORGANIZATION REPORT NUMBER ARCCB-TR-93016	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army ARDEC Close Combat Armaments Center Picatinny Arsenal, NJ 07806-5000			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Published in <i>Physical Review A15</i> (American Physical Society)				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Mandelbrot's rule for sections is generalized to apply to the Hentschel and Procaccia fractal dimension at arbitrary q and on arbitrary sections. It is shown that for almost all $(n-m)$ -dimensional sections, $D^q(q) = D^{n-m}(q) + m$, where the $D^q(q)$ are box-counting Hentschel and Procaccia generalized fractal dimensions of r -dimensional sections of homogeneous fractal point sets in R^n . The rule is shown to apply for finite "thickness" sections as well as "true" sections. A more general form of the rule applicable to inhomogeneous fractal sets is also presented.				
14. SUBJECT TERMS Fractals, Generalized Dimensions, Box-Counting, Fractal Dimensions			15. NUMBER OF PAGES 8	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	ii
INTRODUCTION	1
THEORY	1
SUMMARY AND DISCUSSION	4
REFERENCES	5

List of Illustrations

1. Illustration of the box and strip definitions in R^2	6
---	---

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

ACKNOWLEDGEMENTS

The author gratefully acknowledges P.J. Cote and Mark Johnson of Benet Laboratories for stimulating discussions and encouragement concerning this work.

INTRODUCTION

Mandelbrot's rule (ref 1) states that for almost all $(n-1)$ -dimensional sections

$$D_{HB}^n = D_{HB}^{n-1} + 1$$

where the D_{HB}^r are Hausdorff-Bouligand dimensions of r -dimensional sections of homogeneous fractal sets in R^n . This important rule comprises the basis for a substantial body of numerical fractal analysis.

The following generalization of Mandelbrot's rule is demonstrated here: For almost all $(n-m)$ -dimensional sections

$$D^n(q) = D^{n-m}(q) + m \quad (1)$$

where the $D^r(q)$ are box-counting Hentschel and Procaccia (refs 2,3) generalized fractal dimensions of r -dimensional sections of homogeneous fractal point sets in R^n . Equation (1) obtains for arbitrary q and $n > m \geq 1$. The $q = 0, m = 1$ form of Eq. (1) is the box-counting formulation of the standard Mandelbrot rule.

THEORY

Let the point set in R^n be represented as points in an arbitrarily-oriented Cartesian coordinate system. Define an ensemble of $(n-m)$ -dimensional "sections,"

$$\begin{aligned} T_{n,m}(J) &\equiv T_{n,m}(X_i(J) | i = 1, m; \delta) \\ &\equiv \{ \bar{x} | X_i(J) < x_i \leq X_i(J) + \delta, i = 1, m; -L < x_j \leq L; j = m + 1, n \} \end{aligned} \quad (2)$$

where L is large enough to cover a substantial portion of the fractal set, δ is "small," and the $\{X_i(J)\}$ are such that the $T_{n,m}(J)$ fill a region of R^n without overlaps or gaps. We refer to $T_{n,m}(J)$ sections as "J-strips."

Define sets of boxes (hypercubes) of side E aligned along the coordinate axes in R^n . Let each hypercube in R^n be identified by the position of one of its corners in units of E , i.e., the box at $(I(1), \dots, I(n))E$ is indexed $[I(1), \dots, I(n)]$. Such sets of boxes define corresponding sets of boxes of side E and "thickness" δ^m in each J-strip; the box corresponding to the $[I(1), \dots, I(n)]$ box, which intersects the J-strip, induces the box, which is indexed $[J; I(m+1), \dots, I(n)]$, in the J-strip. There will be approximately E/δ J-strips, which intersect a given box in R^n . Figure 1 illustrates the definitions in R^2 .

Box-counting fractal analysis is generally applied to large, but finite, subsets of fractal sets. The fractal subsets are assumed to be large enough to yield converged box-counting dimensions. Small δ implies that $\delta \ll E_{\min}$, where E_{\min} is the smallest hypercube edge. The present results also apply for complete fractal sets and in the limit $\delta \rightarrow 0$.

Let $p_{I(1), \dots, I(n)}$ denote the occupation probability for the $[I(1), \dots, I(n)]$ box of side E in R^n , which is defined as the number of members of the subset of the fractal point set in the $[I(1), \dots, I(n)]$ box divided by the number of members of the subset of the fractal point set in all the boxes in R^n .

Let $N_{J; I(m+1), \dots, I(n)}$ be the number of points of the fractal subset in the $[J; I(m+1), \dots, I(n)]$ box.

Let S_J be the number of members of the subset of the fractal point set in the J-strip:

$$S_J = \sum_{I(m+1), \dots, I(n)} N_{J; I(m+1), \dots, I(n)} .$$

S_J is proportional to the J-strip probability σ_J in R^n , where

$$\sigma_J \equiv S_J/N$$

$$\text{and } N = \sum_J S_J = \sum_{J; I(m+1), \dots, I(n)} N_{J; I(m+1), \dots, I(n)}$$

is the total number of members in the fractal subset. The occupation probability for the $[J; I(m+1), \dots, I(n)]$ box is $\pi_{J; I(m+1), \dots, I(n)} \equiv N_{J; I(m+1), \dots, I(n)} / S_J$. By definition

$$\sum_{I(m+1), \dots, I(n)} \pi_{J; I(m+1), \dots, I(n)} = 1 .$$

The partition function $Z^n(q, E)$ in R^n is defined as

$$Z^n(q, E) \equiv \sum_{I(1), \dots, I(n)} p_{I(1), \dots, I(n)}^q . \quad (3a)$$

Since the J-strip box corresponding to the $[I(1), \dots, I(n)]$ box in R^n has volume $\Omega_J = E^n (\delta/E)^m$,

$$p_{I(1), \dots, I(n)} \approx (E/\delta)^m \sigma_J \pi_{J; I(m+1), \dots, I(n)} . \quad (3b)$$

Furthermore, each box in R^n is counted approximately $(E/\delta)^m$ times in the sum over strips. Thus, the partition function $Z^n(q, E)$ in R^n can be expanded as

$$\begin{aligned} Z^n(q, E) &= \sum_J \sum_{I(m+1), \dots, I(n)} \left[\frac{\delta}{E} \right]^m \left(\left[\frac{E}{\delta} \right]^m \pi_{J; I(m+1), \dots, I(n)} \sigma_J \right)^q \\ &= \left[\frac{E}{\delta} \right]^{(q-1)m} \sum_J \sigma_J^q Z^{n-m}(J; q, E) . \end{aligned} \quad (3c)$$

where $Z^{n-m}(J; q, E)$ is the J-strip partition function. The approximations become exact in the limit $E \rightarrow 0$ with $E/\delta \gg 1$ and are "good" approximations in the spirit of box-counting algorithms for "small enough" E with $E/\delta \gg 1$; exact equalities obtain in the $E \rightarrow 0$ limit for true sections also.

Box-counting generalized fractal dimensions are defined as

$$D^n(q) = \left\{ \frac{\partial \ln(Z^n(q, E))}{\partial \ln(E)} \right\}_{E_0} \quad (4a)$$

$$D^{n-m}(J; q) = \left\{ \frac{\partial \ln(Z^{n-m}(J; q, E))}{\partial \ln(E)} \right\}_{E_0} \quad (4b)$$

where the right-hand sides are usually determined by least squares fitting over a range of "small" yardstick values centered about a yardstick E_0 .

Since $\ln(Z^n(q, E))$ and $\ln(Z^{n-m}(J; q, E))$ scale like $(q-1)D^n(q)\ln(E)$ and $(q-1)D^{n-m}(J; q)\ln(E)$, respectively, Eq. (4) implies that

$$D^n(q) = \langle D^{n-m}(q) \rangle_{E_0} + m \quad (5)$$

where

$$\langle D^n(q) \rangle_{E_0} = \sum_J D^n(J; q) Z^n(J; q, E_0) \sigma_J^q / \sum_{J'} Z^n(J'; q, E_0) \sigma_{J'}^q. \quad (6)$$

(Since the strip partition functions $Z^{n-m}(J; q, E_0)$ are of the same order of magnitude for given q and E_0 ,

$$\langle D^n(q) \rangle_{E_0} \approx \sum_J D^n(J; q) \sigma_J^q / \sum_{J'} \sigma_{J'}^q, \quad (6')$$

might be a useful approximation.)

N.B., Eqs. (5), (6), and (6') may be interpreted for inhomogeneous fractal sets.

Excluding "exceptional sections," the $D^n(J; q)$ are independent of J for homogeneous fractal point sets and Eq. (5) is equivalent to the generalized form of Mandelbrot's rule for box-counting fractal dimensions at arbitrary q and almost all $(n-m)$ -dimensional J -strips

$$D^n(q) = D^{n-m}(J; q) + m, \text{ for almost all } J. \quad (7)$$

The existence of "exceptional sections" related to special symmetries in fractal point sets, as discussed by Mandelbrot (ref 1), necessitates the inclusion of the "almost all" caveat in the statement of the rule for sections.

The steps in the derivation of Eqs. (5) through (7) hold in the $\delta \rightarrow 0$ limit. Thus, the principal results hold not only for finite "thickness" (finite δ) sections, but they also hold for "true" sections of fractal sets.

SUMMARY AND DISCUSSION

Mandelbrot's rule relates the fractal dimension (for $q = 0$) of a fractal point set in R^n , $D^0(0)$, to the fractal dimension (for $q = 0$) of subsets of the fractal point set in $(n-1)$ -dimensional sections, $D^{n-1}(0)$. The principal results of the present work, contained in Eqs. (5) through (7), generalize Mandelbrot's rule for sections to apply for box-counting Hentschel and Procaccia fractal dimensions $D^q(q)$ at arbitrary q for r -dimensional sections of small "thickness" δ with $n \geq r > 0$. Small δ implies that $\delta \ll E_n$, where E_n is defined after Eq. (4). Furthermore, the equations hold in the $\delta \rightarrow 0$ limit, and thus apply for "true" sections.

One might work with finite δ sections to analyze point sets generated by nonlinear mappings or the distribution of galaxies. One might work with "true" two-dimensional sections of three-dimensional surfaces in a multifractal generalization of the slit-island method (ref 4) (which determines $D(0)$ of fracture surfaces by perimeter-area analysis of fracture surface sections).

Since sparse regions of a fractal subset dominate the average in Eq. (6) for $q < 0$, numerical problems in box-counting evaluations of $D^q(q)$ are likely at $q < 0$. Modified box-counting strategies, which put less stress on sparsely occupied strips, might improve fractal analysis at negative q .

REFERENCES

1. B.B. Mandelbrot, *Fractal Geometry of Nature*, Freeman, New York, 1983.
2. H.G.E. Hentschel and I. Procaccia, *Physica D*, Vol. 8, 1983, p. 435.
3. Thomas C. Halsey, Mogens H. Jensen, Leo P. Kadanoff, Itamar Procaccia, and Boris I. Shraiman, *Physical Review A*, Vol. 33, 1986, p. 1141.
4. B.B. Mandelbrot, D.E. Passoja, and A.J. Paullay, *Nature*, Vol. 308, 1984, p. 721.




Figure 1. Illustration of the box and strip definitions in R^2 . The box indexed $[I(1), I(2)]$ of side E , whose "volume" is E^2 in R^2 , induces the box indexed $[J; I(2)]$ of side E in the J -strip. The J -strip is a one-dimensional section of "thickness" δ .

TECHNICAL REPORT INTERNAL DISTRIBUTION LIST

	NO. OF COPIES
CHIEF, DEVELOPMENT ENGINEERING DIVISION	
ATTN: SMCAR-CCB-DA	1
-DC	1
-DI	1
-DR	1
-DS (SYSTEMS)	1
CHIEF, ENGINEERING SUPPORT DIVISION	
ATTN: SMCAR-CCB-S	1
-SD	1
-SE	1
CHIEF, RESEARCH DIVISION	
ATTN: SMCAR-CCB-R	2
-RA	1
-RE	1
-RM	1
-RP	1
-RT	1
TECHNICAL LIBRARY	5
ATTN: SMCAR-CCB-TL	
TECHNICAL PUBLICATIONS & EDITING SECTION	3
ATTN: SMCAR-CCB-TL	
OPERATIONS DIRECTORATE	1
ATTN: SMCWV-ODP-P	
DIRECTOR, PROCUREMENT DIRECTORATE	1
ATTN: SMCWV-PP	
DIRECTOR, PRODUCT ASSURANCE DIRECTORATE	1
ATTN: SMCWV-QA	

NOTE: PLEASE NOTIFY DIRECTOR, BENET LABORATORIES, ATTN: SMCAR-CCB-TL, OF ANY ADDRESS CHANGES.

TECHNICAL REPORT EXTERNAL DISTRIBUTION LIST

	NO. OF COPIES		NO. OF COPIES
ASST SEC OF THE ARMY RESEARCH AND DEVELOPMENT ATTN: DEPT FOR SCI AND TECH THE PENTAGON WASHINGTON, D.C. 20310-0103	1	COMMANDER ROCK ISLAND ARSENAL ATTN: SMCRI-ENM ROCK ISLAND, IL 61299-5000	1
ADMINISTRATOR DEFENSE TECHNICAL INFO CENTER ATTN: DTIC-FDAC CAMERON STATION ALEXANDRIA, VA 22304-6145	12	MIAC/CINDAS PURDUE UNIVERSITY P.O. BOX 2634 WEST LAFAYETTE, IN 47906	1
COMMANDER US ARMY ARDEC ATTN: SMCAR-AEE	1	COMMANDER US ARMY TANK-AUTMV R&D COMMAND ATTN: AMSTA-DDL (TECH LIB) WARREN, MI 48397-5000	1
SMCAR-AES, BLDG. 321	1	COMMANDER	
SMCAR-AET-O, BLDG. 351N	1	US MILITARY ACADEMY	1
SMCAR-CC	1	ATTN: DEPARTMENT OF MECHANICS	
SMCAR-CCP-A	1	WEST POINT, NY 10996-1792	
SMCAR-FSA	1		
SMCAR-FSM-E	1	US ARMY MISSILE COMMAND	
SMCAR-FSS-D, BLDG. 94	1	REDSTONE SCIENTIFIC INFO CTR	2
SMCAR-IMI-I (STINFO) BLDG. 59	2	ATTN: DOCUMENTS SECT, BLDG. 4484	
PICATINNY ARSENAL, NJ 07806-5000		REDSTONE ARSENAL, AL 35898-5241	
DIRECTOR US ARMY BALLISTIC RESEARCH LABORATORY ATTN: SLCBR-DD-T, BLDG. 305	1	COMMANDER US ARMY FGN SCIENCE AND TECH CTR ATTN: DRXST-SD	1
ABERDEEN PROVING GROUND, MD 21005-5066		220 7TH STREET, N.E. CHARLOTTESVILLE, VA 22901	
DIRECTOR US ARMY MATERIEL SYSTEMS ANALYSIS ACTV ATTN: AMXSY-MP	1	COMMANDER US ARMY LABCOM	
ABERDEEN PROVING GROUND, MD 21005-5071		MATERIALS TECHNOLOGY LAB ATTN: SLCMT-IML (TECH LIB)	2
COMMANDER HQ, AMCCOM		WATERTOWN, MA 02172-0001	
ATTN: AMSMC-IMP-L	1		
ROCK ISLAND, IL 61299-6000			

NOTE: PLEASE NOTIFY COMMANDER, ARMAMENT RESEARCH, DEVELOPMENT, AND ENGINEERING CENTER, US ARMY AMCCOM, ATTN: BENET LABORATORIES, SMCAR-CCB-TL, WATERVLIET, NY 12189-4050, OF ANY ADDRESS CHANGES.

TECHNICAL REPORT EXTERNAL DISTRIBUTION LIST (CONT'D)

	<u>NO. OF COPIES</u>		<u>NO. OF COPIES</u>
COMMANDER US ARMY LABCOM, ISA ATTN: SLCIS-IM-TL 2800 POWDER MILL ROAD ADELPHI, MD 20783-1145	1	COMMANDER AIR FORCE ARMAMENT LABORATORY ATTN: AFATL/MN EGLIN AFB, FL 32542-5434	1
COMMANDER US ARMY RESEARCH OFFICE ATTN: CHIEF, IPO P.O. BOX 12211 RESEARCH TRIANGLE PARK, NC 27709-2211	1	COMMANDER AIR FORCE ARMAMENT LABORATORY ATTN: AFATL/MNF EGLIN AFB, FL 32542-5434	1
DIRECTOR US NAVAL RESEARCH LAB ATTN: MATERIALS SCI & TECH DIVISION CODE 26-27 (DOC LIB) WASHINGTON, D.C. 20375	1 1	DIRECTOR US ARMY BALLISTIC RESEARCH LABORATORY ATTN: SLCBR-IB-M (DR. BRUCE BURNS) ABERDEEN PROVING GROUND, MD 21005-5066	1

NOTE: PLEASE NOTIFY COMMANDER, ARMAMENT RESEARCH, DEVELOPMENT, AND ENGINEERING CENTER, US ARMY AMCCOM, ATTN: BENET LABORATORIES, SMCAR-CCB-TL, WATERVLIET, NY 12189-4050, OF ANY ADDRESS CHANGES.